

MSLC Workshop Series

Math 1151 – Workshop

Sigma Notation and Riemann Sums

Sigma Notation:

Notation and Interpretation of $\sum_{k=1}^n a_k = a_1 + a_2 + a_3 + a_4 + \dots + a_{n-1} + a_n$

- Σ (capital Greek sigma, corresponds to the letter S) indicates that we are to **sum** numbers of the form indicated by the general term
- a_k is the **general term**, which determines what is being summed, and can be defined however we want but is usually a formula containing the index: $a_k = f(k)$
- k is called the **index**; we may use any letter for the index, typically we use $i, j, k, l, m,$ and n as indices
- The index runs through the positive integers, starting with the number below the Σ (in this case 1) and ending with the integer above the Σ (in this case n)
- The sum on the right-hand side is the **expanded form**. (The \dots contains all the terms I was too lazy to write.)
- The letter below the sigma is the variable with respect to the sum. All other letters are **constants** with respect to the sum.

Examples:

1. $\sum_{k=1}^7 k =$

2. $\sum_{k=3}^8 \frac{1}{k+2} =$

3. $\sum_{k=1}^9 4 =$

Special Sum Formulas

$$\sum_{i=1}^n 1 = n$$

$$\sum_{i=1}^n i = \frac{n(n+1)}{2}$$

$$\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

$$\sum_{i=1}^n i^3 = \left[\frac{n(n+1)}{2} \right]^2$$

Examples:

1. $\sum_{k=1}^{200} k =$

2. $\sum_{i=1}^{30} i^2 =$

Properties of Sigma Algebra Σ is an **operator** that represents summation, and its properties are similar to the properties of addition (note what properties are **not** mentioned here)

- Multiplication by a common constant (also called a **scalar** multiple) $\sum c a_k = c \sum a_k$
- Addition or Subtraction (this is also called the **linearity** property) $\sum (a_k \pm b_k) = \sum a_k \pm \sum b_k$

Example:

$$\sum_{k=1}^{10} (2k^2 + k + 1) =$$

Riemann Sums: $R = \sum_k(\text{height of } k\text{th rectangle}) \cdot (\text{width of } k\text{th rectangle})$




FIGURE 5.8

Definition of a Riemann Sum:

Consider a function $f(x)$ defined on a closed interval $[a, b]$, partitioned into n subintervals of equal width by means of grid points $a = x_0 < x_1 < x_2 < \dots < x_{n-1} < x_n = b$. On each subinterval $[x_{k-1}, x_k]$, pick a sample point x_k^* . Then the Riemann sum for f corresponding to this partition is given by:

$$R = \sum_{k=1}^n f(x_k^*)\Delta x = f(x_1^*)\Delta x + f(x_2^*)\Delta x + \dots + f(x_n^*)\Delta x$$

• **WIDTH:** Δx

Since we partition the interval into evenly spaced partitions, we can calculate the width:

$$\Delta x = \frac{b-a}{n}, \text{ where } n \text{ is the number of partitions.}$$

• **HEIGHT:** $f(x_k^*)$

Also, we usually don't pick x_k^* arbitrarily. We use a rule to pick x_k^* . The most common rules to use are the Right Endpoint Rule, the Left Endpoint Rule, and the Midpoint Rule.

The most common rules to use are:

Right Endpoint Rule $x_k^* =$

Left Endpoint Rule $x_k^* =$

Midpoint Rule $x_k^* =$

- If we partition the interval into more and more rectangles with smaller and smaller widths, we get closer to the (signed) area trapped between the curve $y = f(x)$ and the x -axis.

This is where **Sigma Notation** comes in because it becomes time consuming to add up all the terms when there are many, many rectangles.

Calculating A Riemann Sum

Using the **Right Endpoint Rule**, the Riemann sum becomes:

$$\sum_{k=1}^n f\left(a + k\Delta x\right) (\Delta x) = \sum_{k=1}^n \left(\frac{b-a}{n}\right) f\left(a + k\frac{b-a}{n}\right)$$

Using the **Left Endpoint Rule**, the Riemann sum becomes:


$$\sum_{k=1}^n f\left(a + (k-1)\Delta x\right) (\Delta x) = \sum_{k=1}^n \left(\frac{b-a}{n}\right) f\left(a + (k-1)\frac{b-a}{n}\right)$$

Using the **Midpoint Rule**, the Riemann sum becomes:

$$\sum_{k=1}^n f\left(a + \left(\frac{(k-1) + k}{2}\right)\Delta x\right) (\Delta x) = \sum_{k=1}^n \left(\frac{b-a}{n}\right) f\left(a + \left(\frac{(k-1) + k}{2}\right)\frac{b-a}{n}\right)$$

Example:


Estimate the area under $f(x) = x^2 + 2$ on the interval $[-2, 3]$ using right Riemann Sums and 5 rectangles.



NOVICE (before Calculus):

Example:


Estimate the area under $f(x) = x^2 + 2$ on the interval $[-2, 3]$ using right Riemann Sums and 5 rectangles.



SEMI-PRO (beginning Calculus student):


Example:

Estimate the area under $f(x) = x^2 + 2$ on the interval $[-2, 3]$ using right Riemann Sums and 5 rectangles.




PRO (by test time):

Left Sum: Estimate the area under $f(x) = x^2 + 2$ on the interval $[-2, 3]$ using left Riemann Sums and 5 rectangles.



Midpoint Sum: Estimate the area under $f(x) = x^2 + 2$ on the interval $[-2, 3]$ using midpoint Riemann Sums and 5 rectangles.



Being More Accurate:

What if we want to get a better approximation than any of the above give us? More rectangles will give us less extra or unused area between the curve and the x -axis. Let's do a **right sum** for $\int_{-2}^3 (x^2 + 2) dx$ using 5000 equal subintervals.

First calculate the width: $\Delta x =$

Then the x -value for the right endpoint of the k th rectangle is:

Thus the height of the k th rectangle is:

So the Riemann sum is

Now evaluate this sum using your knowledge of sigma algebra!